

Patient  
Information  
for Consent

RESTORE  
PLASTIC SURGERY

CM04 Abdominoplasty

Expires end of March 2024

Augustine Akali - Consultant Plastic Surgeon  
01482 223 553  
07487 584 911  
practice.manager@restoresurgery.com  
www.restoresurgery.com

Get more information and references at [www.aboutmyhealth.org](http://www.aboutmyhealth.org)  
Tell us how useful you found this document at [www.patientfeedback.org](http://www.patientfeedback.org)



Royal College  
of Surgeons  
of England



THE ROYAL  
COLLEGE OF  
SURGEONS  
OF EDINBURGH



Patient Information Forum

[eidohealthcare.com](http://eidohealthcare.com)

EIDO  
HEALTHCARE

UNITED KINGDOM

## What is an abdominoplasty?

An abdominoplasty (or 'tummy tuck') is an operation to improve the appearance of your abdomen. It may involve removing excess skin and fat and tightening your abdominal muscles.

Your surgeon will assess you and tell you if an abdominoplasty is suitable for you. However, it is your decision to go ahead with the operation or not.

This document will give you information about the benefits and risks to help you to make an informed decision. If you have any questions that this document does not answer, it is important that you ask your surgeon.

Once all your questions have been answered and you feel ready to go ahead with the procedure, you will be asked to sign the informed consent form. However, you can still change your mind at any point.

## Is an abdominoplasty suitable for me?

It is common for the skin of your abdomen not to shrink back after pregnancy or losing a lot of weight. You are most likely to benefit from an abdominoplasty if your abdomen makes you feel uncomfortable or you are self-conscious about what your abdomen looks like.

Your surgeon will carry out a detailed assessment before deciding if surgery is suitable for you. This may include taking photos for your medical records. They will examine your abdomen and ask you questions about your medical history.

Your surgeon will check if you are the right weight for your height. If you are overweight, it is better to lose weight before having surgery.

Let your surgeon know if you are pregnant or planning to become pregnant in the future. Pregnancy can change the appearance of your abdomen and may affect the long-term results of surgery.

## What are the benefits of surgery?

Your abdomen should be flatter and have a better appearance. If you have a gap between your abdominal muscles causing you to feel uncomfortable, this will be improved after surgery.

Most people who have a successful abdominoplasty are more comfortable with their appearance, are able to wear better fitting clothing and their personal and sexual relationships may improve.

## Are there any alternatives to an abdominoplasty?

If your abdomen looks saggy because the muscles of your abdominal wall are slack, your appearance may improve if you follow an exercise programme.


If the problem is caused mainly by too much fat, it may be better to have only liposuction, where the fat is sucked out using a small tube or the fat is frozen.

Your surgeon may recommend having liposuction before, during or after an abdominoplasty. Liposuction is an essential part of certain types of abdominoplasty.


## What does the operation involve?

The healthcare team will carry out a number of checks to make sure you have the operation you came in for. You can help by confirming to your surgeon and the healthcare team your name and the operation you are having.

The operation is performed under a general anaesthetic and usually takes 2 to 5 hours. You may also have injections of local anaesthetic to help with the pain after the operation. You may be given antibiotics during the operation to reduce the risk of infection.



A standard abdominoplasty



The results of a standard abdominoplasty

fat between your pubic area and your umbilicus (belly button).

Your surgeon will cut under the skin above your umbilicus then pull it downwards, tightening it.

Your surgeon will need to make a small cut for your umbilicus to be positioned as close to the middle of your tummy as possible. Sometimes they will need to reconstruct your umbilicus (umbilicoplasty).

Your surgeon may need to tighten the muscles of your abdominal wall and use liposuction to remove excess fat.

### Fleur-de-lis abdominoplasty

This technique is suitable if you have a lot of excess skin hanging downwards and around your abdomen, usually as a result of losing a significant amount of weight or having bariatric surgery. The procedure is similar to a standard abdominoplasty. Your surgeon will also tighten your waist by making a vertical cut along your midline up to just below your breast bone, to remove more excess skin and fat.

### Mini-abdominoplasty

Your surgeon will need to make a smaller cut just above the pubic area to remove the excess skin. A smaller amount of skin is removed during a mini-abdominoplasty than with a full abdominoplasty.

Your surgeon may need to tighten the muscles of your abdominal wall. They will not need to cut under the skin above your umbilicus and will use liposuction only if they need to remove any excess fat in the upper part of your tummy.

### Lower bodylift or belt lipectomy

These techniques are suitable after significant weight loss, usually as a result of bariatric surgery.


Your surgeon will remove excess skin and fat from your sides, abdomen and lower back. They may also use liposuction to remove excess fat from your thighs.

The following are the main techniques that your surgeon may use and they will discuss with you which technique is most appropriate for you.


### Standard abdominoplasty

This technique is suitable if you have a lot of excess skin.

Your surgeon will make a curved cut from one hip to the other, going down to just above your pubic area. They will remove any excess skin and



The abdominal muscles being stitched together



A mini-abdominoplasty

### Endoscopic abdominoplasty

This technique is not commonly used as it is only suitable if you need to have the muscles of your abdominal wall tightened and your skin has lost a lot of its elasticity. That is uncommon unless you have a hernia around your belly button. If you want more information speak to your surgeon.

## What should I do about my medication?

Make sure your healthcare team knows about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.

## How can I prepare myself for the operation?

If you smoke or use nicotine products, stopping smoking at least 6 weeks before and for 6 weeks after surgery may reduce your risk of developing complications and will improve your long-term health. Many surgeons will not perform an abdominoplasty if the patient is still smoking or using nicotine.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound.

- In the week before the operation, do not shave or wax the area where a cut is likely to be made.
- Have a bath or shower either the day before or on the day of the operation.
- Keep warm around the time of the operation. Let the healthcare team know if you feel cold.
- If you are diabetic, keep your blood sugar levels under control around the time of your procedure. Your surgeon will discuss the increased risk of complications if you are a diabetic.

Speak to the healthcare team about any vaccinations you might need to reduce your risk of serious illness while you recover. When you come into hospital, practise social distancing and

hand washing and wear a face covering when asked.

## What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you. Some risks are higher if you are older, obese, you are a smoker or have other health problems. These health problems include diabetes, heart disease or lung disease.

Some complications can be serious and can even cause death.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

## General complications of any operation

- Bleeding during or soon after the operation. Rarely, you will need a blood transfusion or another operation. It is important you are not on blood thinners at the time of the operation.
- Infection of the surgical site (wound). It is usually safe to shower after 2 days but you should check with the healthcare team. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. In a standard abdominoplasty minor infections are common because the central part of your wound is often slow to heal. Any serious infection usually needs treatment with antibiotics or another operation and can make the scar more noticeable. Do not take antibiotics unless you are told you need them. If you have a high fever and severe pain in your stomach, you must call the Emergency department.
- Blood clot in your leg (deep-vein thrombosis – DVT). This can cause pain, swelling or redness in your leg, or the veins near the surface of your leg to appear larger than

normal. The healthcare team will assess your risk. They will encourage you to get out of bed soon after the operation and may give you injections, medication, or special stockings to wear. Special footwear that compresses your leg and increases blood flow will be used when you are asleep and until you are mobile. Let the healthcare team know straight away if you think you might have a DVT once you are at home. A DVT may cause pain and swelling in your legs, usually starting a few days after the operation.

- Blood clot in your lung (pulmonary embolus), where a blood clot moves through your bloodstream to your lungs. Let the healthcare team know straight away if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. This usually happens once you are at home. If this happens, call an ambulance or go immediately to your nearest Emergency department.
- Allergic reaction to the equipment, materials or medication. The healthcare team is trained to detect and treat any reactions that might happen. Let the healthcare team know if you have any allergies or if you have reacted to any medication or tests in the past.
- Chest infection. Your risk will be lower if you have stopped smoking and you are free of Covid-19 (coronavirus) symptoms for at least 7 weeks before the operation.

## Specific complications of this operation

- Developing a collection of blood (haematoma) or fluid (seroma) under the skin in your lower abdomen. You may need another operation to remove the blood or fluid. After that you may need a blood transfusion.
- Numbness or continued pain on your abdomen caused by injury to the small nerves that supply your skin. Any pain or numbness usually gets better within a few weeks but can sometimes continue for many months.
- Wound-healing problems, if some areas do not heal properly or some skin dies. You may need to come back to the clinic often to have the dressings changed. Occasionally you may



need further surgery to remove some of the tissue.

- **Cosmetic problems.** It is difficult to predict exactly how your abdomen will appear after the operation. Sometimes you will still have a small area of excess skin. You may get flaps of skin towards your sides near the ends of your wound. If you had an abdominoplasty, your umbilicus may go slightly off-centre but usually only you will notice this. If you had a fleur-de-lis abdominoplasty, your midline scar may thicken and become more noticeable. It is not possible to guarantee the results of the operation.

## Consequences of this procedure

- Pain is usually only mild and easily controlled with simple painkillers such as paracetamol. However, you can feel sore if you had a standard abdominoplasty or if your surgeon had to tighten your abdominal muscles.
- Unsightly scarring of your skin. The scars usually settle with time. If you have dark skin, the scars can sometimes stay thick and red. The scar from the main cut will be noticeable but you will usually be able to hide most of the scar, even when wearing underwear or a swimming costume. If you have a standard abdominoplasty you will also have a scar around your umbilicus. Follow the instructions your surgeon gives you about how to care for your wounds. Your surgeon will discuss scarring with you.

## How soon will I recover?

### In hospital

After the operation you will be transferred to the recovery area and then to the ward. Your abdomen will look discoloured and feel firm and swollen.

Try to keep your knees bent to reduce the tension on your stitches. Ask for some pillows to be put behind your knees.

You should be able to go home after 1 to 3 days. However, your doctor may recommend that you stay a little longer depending on your progress.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

## Returning to normal activities

To reduce the risk of a blood clot, make sure you carefully follow the instructions of the healthcare team if you have been given medication or need to wear special stockings.

Rest for 2 weeks and do only light activities. This will help your wound to heal and reduce the risk of fluid collecting. You should then be able to do a limited amount of activity, such as lifting young children.

You should be able to return to work after 2 weeks, depending on your type of work.

Do not lift anything heavy or do strenuous exercise, such as vacuuming or ironing, for 6 weeks.

Do not have sex until your stomach is no longer sore and then be gentle.

The healthcare team will advise you on exercising to help keep you to a healthy weight and to improve the tone of your abdominal muscles.

Do not drive until you can control your vehicle, including in an emergency, and always check your insurance policy and with the healthcare team.

## The future

The healthcare team will arrange for you to come back to the clinic regularly to check on your progress.

The results of an abdominoplasty improve gradually over the first 6 months. The best results for you will happen if you keep to a healthy weight and exercise regularly.

If you put on a lot of weight and then lose weight, or become pregnant, you may get too much skin on your abdomen again. However, you should not get as much excess skin as you would have done without the operation.

## Summary

An abdominoplasty is an operation to improve the appearance of your abdomen. You should

consider the options carefully and have realistic expectations about the results.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

## **Acknowledgements**

### **Reviewer**

Eoin O'Broin (MD, FRCS (Plast.))

### **Illustrator**

Medical Illustration Copyright © Nucleus Medical Art. All rights reserved. [www.nucleusinc.com](http://www.nucleusinc.com)